

8

LITERATURE CITED

- Aarssen, L. W. (1997), High productivity in grassland systems: affected by species diversity or productive species? *OIKOS*. 87:183-184.
- Aarssen, L. W. (2001), On correlations and causations between productivity and species richness: predictions from habitat attributes. *Basic and Applied Ecology*. 2:105-114.
- Anderson, R. C. (1990), *The historic role of fire in the North American grassland*. In *Effects of fire on Tallgrass Prairie Ecosystems* (S. L. Collins, and L. Wallace, Eds.), Oklahoma University Press, Norman, OK, USA.
- Asner, G. P. (1998), Biophysical and biochemical sources of variability in canopy reflectance. *Remote Sensing of Environment*. 64:234-253.
- Asner, G. P., Wessman, C. A., Schimel, D. S., and Archer, S. (1998), Variability in leaf and litter optical properties: implications for BRDF model inversions using AVHRR, MODIS and MISR. *Remote Sensing of Environment*. 63:243-257.
- Asrar, G., Weiser, R. L., Johnson, D. E., Kanemasu, E. T., and Milleen, J. M. (1986), Distinguishing among tallgrass prairie cover types from measurements of multispectral reflectance. *Remote Sensing of Environment*. 19:159-169.
- Atkinson, P. M. (1999), Geographic Information Science: Geostatistics and Uncertainty. *Progress in Physical Geography*. 23(1):134-142.
- Axelrod, D. I. (1985), Rise of the grassland biome, Central North America. *The Botanical Review*. 51:163-201.
- Bahr, L. M. (1971), Nutritional strategies of myomorph rodents in North America. *Journal of Mammology*. 52:800-805.
- Bailey, T. C., and Gatrell, A. C. (1995), *Interactive Spatial Data Analysis*. Longman Scientific and Technical, Essex, UK, pp. 413.
- Baret, F., and Guyot, G. (1991), Potentials and limits of vegetation indices for LAI and APAR assessment. *Remote Sensing of Environment*. 35:161-173.
- Barkman, J. J. (1988), *New systems of plant growth forms and phenological plant types*. In *Plant form and vegetation structure. Adaptation, plasticity and relation to herbivory* (M. J. A. Werger et al., Eds), SPB Academic Publishing, The Hague. pp 9-44.
- Barnes, P. W., Tieszen, L. L., and Ode, D. J. (1983), Distribution, production and diversity of C3- and C4-dominated communities in a mixed prairie. *Canadian Journal of Botany*. 61:741-751.

- Barrett, E. C., and Curtis, L. F. (1982), *Introduction to Environmental Remote Sensing*. Chapman and Hall, UK, pp. 339.
- Bartha, S., Collins, S. L., Glenn, S. M., and Kertész, M. (1995), Fine-scale spatial organization of tallgrass prairie vegetation along a topographic gradient. *Folia Geobot. Phytotax.* 30:169-184.
- Bednekoff, P. A. (2001), Baseline theory of biodiversity and ecosystem function. *Bulletin of the Ecological Society of America.* 82(3):205.
- Belleheumer, C., and Legendre, P. (1998), Multiscale sources of variation in ecological variables: modeling spatial dispersion, elaborating sampling designs. *Landscape Ecology.* 13:15-25.
- Bian, L. (1997), *Multiscale nature of spatial data in scaling up environmental models*. In *Scale in Remote Sensing and GIS* (D. A. Quattrochi, and M. F. Goodchild, Eds.), CRC Press, Boca Raton, FL, USA. pp 13-26.
- Bormann, F. H., and Likens, G. E. (1979), *Pattern and process in a forested ecosystem*. Springer-Verlag, New York, NY, USA, pp. .
- Botkin, D. B. (1975), *Functional groups of organisms in model ecosystems*. In *Ecosystem analysis and prediction* (S. A. Levin, Ed, Society for Industrial and Applied Mathematics, Philadelphia, PA, USA. pp 98-102.
- Bourlière, F., and Hadley, M. (1970), The ecology of tropical savannas. *Annual Review of Ecology and Systematics.* 1:125-152.
- Boutton, T. W., and Tieszen, L. L. (1983), Estimation of plant biomass by spectral reflectance in an east African grassland. *Range Management.* 36(2):213-221.
- Bowman, W. D., and Turner, L. (1993), Photosynthetic sensitivity to temperature in populations of two C4 *bouteloua* (poacea) species native to different altitudes. *American Journal of Botany.* 80(4):369-374.
- Box, E. O., Holben, B. N., and Kalb, H. (1989), Accuracy of the AVHRR vegetation index as a predictor of biomass, primary productivity and net CO2 flux. *Vegetatio.* 80:71-89.
- Brady, N. C. (1990), *The Nature and Properties of Soils*. Collier MacMillan Inc, New York, NY, USA, pp. 621.
- Bragg, T. B. (1995), *The Physical Environment of Great Plains Grasslands*. In *The Changing Prairie. North American Grasslands* (A. Joern, and K. H. Keeler, Eds.), Oxford University Press, New York, NY, USA. pp 49-81.
- Bringmark, E., and Bringmark, L. (1998), Improved soil monitoring by use of spatial patterns. *Ambio.* 27(1):45-52.
- Brown, L. (1989), *The Audubon Society Nature Guides. Grasslands*. Chanticleer Press, Inc, New York, NY, USA, pp. 595.
- Bryson, R. A., and Hare, F. K. (1974), *The Climates of North America*. Climates of North America. Elsevier Scientific Publishing Company, Amsterdam, Holland, pp. 420.

- Bulla, L. (1996), *Relationships between biotic diversity and primary productivity in savanna grasslands*. In *Biodiversity and savanna ecosystem processes: a global perspective* (O. T. Solbrig, E. Medina, and J. F. Silva, Eds.), Springer, New York, NY, USA.
- Burden, R. L., and Faires, J. D. (1997), *Numerical Analysis*. Brooks/Cole Publishing Company, New York, NY, USA, pp. 811.
- Burke, I. C., Kittel, T. G. F., Lauenroth, W. K., Snook, P., Yonker, C. M., and Parton, W. J. (1991), Regional analysis of the central Great Plains. *BioScience*. 41:685-692.
- Burrough, P. A., and McDonnell, R. A. (1998), *Principles of Geographical Information Systems. Spatial Information Systems and Geostatistics*. . Oxford University Press, Oxford, UK., pp. 333 .
- Cain, S. A. (1950), Lifeforms and phytoclimate. *Botanical Review*. 16:1-32.
- Caswell, H., and Cohen, J. E. (1991), *Communities in patchy environments : A model of disturbance, competition and heterogeneity*. In *Ecological Heterogeneity* (J. Kolasa, and S. T. A. Pickett, Eds.), Springer-Verlag, New York, NY, USA. pp 97-123.
- Chapin, F. S. I., Zavaleta, E. S., Eviners, V. T., Naylor, R. L., Vitousek, P. M., Reynolds, H. L., Hooper, D. U., Lavorel, S., Sala, O. E., Hobbie, S. E., Mack, M. C., and Diaz, S. (2000), Consequences of changing biodiversity. *Nature*. 405(11 May 2000):234-242.
- Chen, J. M. (1996), Evaluation of vegetation indices and a modified simple ratio for boreal applications. *Canadian Journal of Remote Sensing*. 22(3):229-242.
- Congalton, R. G. (1991), A review of assessing the accuracy of classifications of remotely sensed data. *Remote Sensing of Environment*. 37:35-46.
- Coppedge, B. R., Engle, D. M., Toepfler, C. S., and Shaw, J. H., (1998), Effects of seasonal fire-bison grazing and climatic variation on tallgrass prairie vegetation. *Plant Ecology*. 139:235-246.
- Coupland, R. T. (1950), Ecology of Mixed Prairie in Canada. *Ecological Monographs*. 20(4):271-315.
- Coupland, R. T. (1961), A reconsideration of grassland classification in the Great Plains of North America. *Journal of Ecology*. 49:135-167.
- Coupland, R. T. (1979), *Grassland Ecosystems of the World*. Cambridge University Press, Cambridge, UK, pp. 401.
- Coupland, R. T. (1992), *Mixed Prairie*. In *Ecosystems of the World. 8A. Natural Grasslands. Introduction and Western Hemisphere*. , Elsevier, Amsterdam, Holland. pp151-179.
- Coupland, R. T., and Rowe, J. S. (1969), *Natural Vegetation of Saskatchewan*. In *Atlas of Saskatchewan* (J. M. Richards, and K. I. Fung, Eds.), University of Saskatchewan, Saskatoon. pp 236.
- Crawley, M. J., Brown, S. L., Heard, M. S., and Edwards, G. R. (1999), Invasion-resistance in experimental grassland communities: species richness or species identity? *Ecology Letters*. 2:140-148.
- Cressie, N. A. (1991), *Statistics for spatial data*. John Wiley and Sons, New York, NY, USA, pp. 900.

- Cressie, N. A. (1993), *Statistics for spatial data*. John Wiley and Sons, New York, NY, USA, pp. 900.
- Cressie, N. A. C. (1985), Fitting variogram models by weighted least squares. *Mathematical Geology*. 17:563-586.
- Cressie, N. A. C., and Hawkins, D. M. (1980), Robust Estimation of the Variogram: I. *Mathematical Geology*. 12:115-125.
- Crippen, R. E. (1990), Calculating the vegetation index faster. *Remote Sensing of Environment*. 34:71-73.
- Crist, E. P., and Cicone, R. C. (1984), Application of the tasseled cap concept to simulated thematic mapper data. *Photogrammetric Engineering and Remote Sensing*. 50:343-352.
- Csillag, and Kabos. (1996), Hierarchical decomposition of variance with applications in environmental mapping based on satellite images. *Mathematical Geology*. 4(28):385-405.
- Csillag, F., Boots, B., Fortin, M.-J., Lowell, K., and Potvin, F. (Unpublished Manuscript), Multiscale characterization of boundaries and landscape ecological patterns.
- Csillag, F., Davidson, A., and Mitchell, F., 1997. Spatio-temporal patterns of productivity and diversity in Grasslands National Park: Scaling remotely sensed data, The Ecological Monitoring and Assessment Network's (EMAN) Third Science Meeting, Saskatoon, Canada.
- Csillag, F., Davidson, A., Peat, H., Wedin, D., Wylie, B., Fargey, P., and Kertész, M., 1996. Spatiotemporal variability : A component in estimating ecological parameters by remote sensing in Grasslands National Park, Canadian Association of Geographers (CAG), Saskatoon, Canada.
- Csillag, F., Kertész, M., Davidson, A., and Mitchell, S. (2001), On the measurement of diversity-productivity relationships in a northern mixed grass prairie (Grasslands National Park, Saskatchewan, Canada). *Community Ecology*. 2(2):145-159.
- Cuevas, E., Brown, S., and Lugo, A. E. (1991), Above- and belowground organic matter storage and production in a tropical pine plantation and a paired broadleaf secondary forest. *Plant Soil*. 135:264-276.
- Cullinan, V. I., Simmons, M. A., and Thomas, J. T. (1997), A bayesian test of hierarchy theory: scaling up variability in plant cover from field to remotely sensed data. *Landscape Ecology*. 12:273-285.
- Cummins, K. W. (1974), Structure and function of stream ecosystems. *BioScience*. 24:631-641.
- Curran, P. J., Dungan, J. L., Macler, B. A., Plummer, S. E., and Peterson, D. L. (1992), Reflectance spectroscopy of fresh whole leaves for the estimation of chemical concentration. *Remote Sensing of Environment*. 39:153-166.
- Czárán, T., and Bartha, S. (1989), The effect of spatial pattern on community dynamics: a comparison of simulated and field data. *Vegetatio*. 83:229-239.
- Czárán, T., and Bartha, S. (1992), Spatiotemporal dynamic models of plant populations and communities. *Trends in Ecological Evolution*. 7:38-42.
- Dahlman, R. C. (1993), CO₂ and plants : revisited. *Vegetatio*. 104/105:339-355.

- Davidson, A., and Csillag, F. (2001), The influence of vegetation index and spatial resolution on a two-date remote sensing derived relation to C4 species coverage. *Remote Sensing of Environment*. 75(1):138-151.
- Davis, F. W., Quattrochi, D. A., Ridd, M. K., Lam, S.-N., Walsh, S. J., Michaelson, J. C., Franklin, J., Stowe, D. A., Johannsen, C. J., and Johnston, C. A. (1991), Environmental analysis using integrated GIS and remotely sensed data: some research needs and priorities. *Photogrammetric Engineering and Remote Sensing*. 57(6):689-697.
- Davis, S. D., and Mooney, H. A. (1986), Water use patterns of four occurring chaparral shrubs. *Oecologia*. 70:172-177.
- Deutsch, C. V., and Journel, A. G. (1992), *GSLIB: Geostatistical Software Library and User's Guide*. . Oxford University Press, New York, NY, USA, pp. 369.
- Dodd, M. B., and Lauenroth, W. K. (1997), The influence of soil texture on the soil water dynamics and vegetation structure of a shortgrass steppe ecosystem. *Plant Ecology*. 84:177-183.
- Dodd, M. B., Lauenroth, W. K., and Welker, J. M. (1998), Differential water resource use by herbaceous and woody plant life-forms in a shortgrass steppe community. *Oecologia*. 117:404-512.
- Duccy, M. J., Moser, W. K., and Ashton, P. M. S. (1996), Effect of fire intensity on understory composition and diversity in a Kalmia-dominated Oak forest, New England, USA. *Vegetatio*. 123:81-90.
- Dungan, J. L. (1999), *Conditional simulation: An alternative to estimation for achieving mapping objectives*. In *Spatial statistics for remote sensing* (A. Stein, Ed), Kluwer Academic, Netherlands. pp 135-152.
- Efron, B., and Tibshirani, R. (1991), Statistical data analysis in the computer age. *Science*. 253:390-395.
- Efron, B., and Tibshirani, R. J. (1993), *An Introduction to the Bootstrap*. Monographs on Statistics and Applied Probability, 57. Chapman and Hall, New York, NY, USA., pp. 436 .
- Ehleringer, J. R., and Field, C. B. (1993), *Scaling ecophysiological processes. Leaf to Globe*. Academic Press, San Diego, CA, USA. pp. 388.
- Ehleringer, J. R., and Monson, K. R. (1993), Evolutionary and ecological aspects of photosynthetic pathway variation. *Annual Review of Ecology and Systematics*. 24:411-439.
- Ehleringer, J. R., Sage, R. F., Flanagan, L. B., and Pearcy, R. W. (1991), Climate change and the evolution of C4 photosynthesis. *Trends in Ecology and Evolution*. 6(3):95-99.
- Ehrlich, P. R., and Ehrlich, A. H. (1981), *Extinction, the causes and consequences of the disappearance of species*. . Random House, New York, NY, USA, pp. 305.
- Elton, C. S. (1958), *The ecology of invasions by animals and plants*. . Methuen, London, UK, pp. 181.
- Englund, E. (1993), *Spatial simulations: environmental applications*. In *Environmental modeling with GIS* (M. F. Goodchild, B. O. Parks, and L. T. Steyaert, Eds.), Oxford University Press, Oxford, UK. pp 432-437.
- Environment Canada (1998), *Canadian Climate Normals 1961-1990. Swift Current A, Saskatchewan*.

- Epstein, H. E., Lauenroth, W. K., Burke, I. C., and Coffin, D. P. (1997), Productivity patterns of C3 and C4 functional types in the US Great Plains. *Ecology*. 78(3):722-731.
- Epstein, H. E., Lauenroth, W. K., Burke, I. C., and Coffin, D. P. (1998), Regional productivities of plant species in the Great Plains of the United States. *Plant Ecology*. 134:173-195.
- Fonseca, C. R., and Ganade, G. (2001), Species functional redundancy, random extinctions and the stability of ecosystems. *Journal of Ecology*. 89:118-125.
- Foody, G. M., and Curran, P. J. (1994), *Scale and environmental remote sensing*. In *Environmental remote sensing from regional to global scales* (G. M. Foody, and P. J. Curran, Eds.), John Wiley and Sons, New York, NY, USA.
- Fortin, M.-J., and Jacquez, G. M. (2000), Randomization tests and spatially autocorrelated data. *Bulletin of the Ecological Society of America*. 81(3):201-205.
- Fortin, M.-J. (1999), *Spatial Statistics in Landscape Ecology*. In *Landscape Ecological Analysis: Issues and Applications* (J. H. Klopatek, and R. H. Gardner, Eds.), Springer Verlag, New York, NY, USA. pp 253-279.
- Fourty, T., Baret, F., Jacquemoud, S., Schmuck, G., and Verdebout, J. (1996), Leaf optical properties with explicit description of its biochemical composition: direct and inverse problems. *Remote Sensing of Environment*. 56:104-117.
- Frank, D. A., and McNaughton, S. J. (1991), Stability increases with diversity in plant communities: empirical evidence from the 1988 Yellowstone drought. *OIKOS*. 62:360-362.
- Freund, R. J., and Little, R. C. (1991), *SAS system for regression*. . SAS Institute, Cary, NC, USA, pp. .
- Friedel, M. H., Bastin, G. N., and Griffin, G. F. (1988), Range assessment and monitoring in arid lands: the derivation of functional groups to simplify vegetation data. *Journal of Environmental Management*. 27:85-97.
- Friedl, M. A., Michaelsen, J., Davis, F. W., Walker, H., and Schimel, D. S. (1994), Estimating grassland biomass and leaf area index using ground and satellite data. *Remote Sensing of Environment*. 15(7):1401-1420.
- Garnier, E., Navas, M.-L., Austin, M. P., Lilley, J. M., and Gifford, R. M. (1997), A problem for biodiversity-productivity studies: How to compare the productivity of multispecific plant mixtures to that of monocultures? *Acta Oecologica*. 18(6):657-670.
- Girard, C. M. (1982), Estimation of phenological stages and physiological stages of grasslands from remote sensing data. *Vegetatio*. 48:219-226.
- Gitay, H., and Noble, I. R. (1997), *What are functional types and how should we seek them?* In *Plant Functional Types. Their relevance to ecosystem properties and global change* (T. M. Smith, H. H. Shugart, and F. I. Woodward, Eds.), Cambridge University Press, Cambridge, UK. pp 3-19.
- Givnish, T. J. (1994), Does diversity beget stability? *Nature*. 371(6493):113-114.

- Goel, N. S. (1988), Models of vegetation canopy reflectance and their use in estimation of biophysical parameters from reflectance data. *Remote Sensing of Environment*. 4:1-212.
- Goetz, S. J. (1997), Multi-sensor analysis of NDVI, surface temperature and biophysical variables at a mixed grassland site. *International Journal of Remote Sensing*. 18(1):71-94.
- Golluscio, R. A., and Sala, O. E. (1993), Plant functional types and ecological strategies in Patagonian forbs. *Journal of Vegetation Science*. 4:839-846.
- Goodchild, M. F., and Quattrochi, D. A. (1997), *Introduction: Scale, multiscaling, remote sensing and GIS*. In *Scale in Remote Sensing and GIS* (D. A. Quattrochi, and M. F. Goodchild, Eds.), CRC Press Inc., Boca Raton, FL, USA. pp 1-12.
- Goodin, D. G., and Henebry, G. M. (1997), A technique for monitoring ecological disturbance in tallgrass prairie using seasonal NDVI trajectories and a discriminant function mixture model. *Remote Sensing of Environment*. 61:270-278.
- Goodin, D. G., and Henebry, G. M. (1998), Seasonality of finely-resolved spatial structure of NDVI and its component reflectances in tallgrass prairie. *International Journal of Remote Sensing*. 19(16):3213-3220.
- Gough, L., Grace, J., and Taylor, K. (1994), The relationship between species richness and community biomass: the importance of environmental variables. *OIKOS*. 70:271-279.
- Goutte, C. (1997), Note on free lunches and cross-validation. *Neural Computation*. 9:1211-1215.
- Graniero, P. A., 1999. The effect of spatiotemporal sampling strategies and data acquisition accuracy on the characterization of a dynamic ecological system. In: R. N. Dave, and T. Sudkamp (Editors), 18th International Conference of the North American Fuzzy Information Processing Society - NAFIPS, New York, New York, USA, pp. 829-833.
- Greig-Smith, P. (1952), The use of random and contiguous quadrats in the study of the structure of plant communities. *Annual Botanical New Series*. 16:293-316.
- Grime, J. P. (1997), Biodiversity and ecosystem function: The debate deepens. *Science*. 277:1260-1261.
- Grime, J. P., Hodgson, J. G., and Hunt, R. (1988), *Comparative plant ecology: A functional approach to common British species*. . Unwin Hyman, London, UK, pp. 742.
- Gustafson, E. J. (1998), Quantifying landscape spatial pattern: What is the state of the art? *Ecosystems*. 1:143-156.
- Guterman, L. (2000), Have Ecologists oversold biodiversity? *The Chronicle of Higher Education*. (October 13 2000):24-26.
- Harper, J. L. (1977), *Population Biology of Plants*. . Academic Press, London, UK, pp. 892.
- Hart, M. M., Reader, R. J., and Klironomos, J. N. (2001), Biodiversity and ecosystem function: Alternative hypotheses or a single theory. *Bulletin of the Ecological Society of America*. 82(1):88-90.
- Hatfield, J. L., Asrar, G., and Kanemasu, E. T. (1984), Intercepted photosynthetically active radiation estimated by spectral reflectance. *Remote Sensing of Environment*. 14:65-75.

- Hawkworth, D. L., and Kalin-Arroyo, M. T. (1995), *Magnitude and distribution of biodiversity*. In *Global Biodiversity Assessment* (V. H. Heywood, Ed, Cambridge University Press, Cambridge, UK. pp 107-192.
- Hector, A. (1998), The effect of diversity on productivity: detecting the role of species complementarity. *OIKOS*. 82:597-599.
- Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M. C., Dimer, M., Dimitrakopoulos, P. G., Finn, J. A., Freitas, H., Giller, P. S., Good, J., Harris, R., Högborg, P., Huss-Danell, K., Joshi, J., Jumpponen, A., Körner, C., Leadley, P. W., Loreau, M., Minns, A., Mulder, C. P. H., O'Donovan, G., Otway, S., Pereira, J. S., Prinz, A., Read, D. J., Scherer-Lorenzen, M., Schulze, E.-D., Siamantziouras, A.-S. D., Spehn, E. M., Terry, A. C., Troumbis, A. Y., Woodward, F. I., Yachi, S., and Lawton, J. H. (1999), Plant diversity and productivity experiments in European grasslands. *Science*. 286:1123-1127.
- Henebry, G. M. (1993), Detecting change in grasslands using measures of spatial dependence with Landsat TM data. *Remote Sensing of Environment*. 46:233-234.
- Henebry, G. M., and Su, H. (1993), Using landscape trajectories to assess the effects of radiometric rectification. *International Journal of Remote Sensing*. 14(12):2417-2423.
- Herben, T., Krahulec, F., Hadincova, F., and Kovarova, M. (1993a), Small-scale spatial dynamics of plant species in a grassland community over six years. *Journal of Vegetation Science*. 4:171-178.
- Herben, T., Krahulec, F., Hadincova, F., and Skalova, H. (1993b), Small-scale variability as a mechanism for large-scale stability in mountain grasslands. *Journal of Vegetation Science*. 4:163-170.
- Hobbs, R. J. (1992), *Function of biodiversity in mediterranean ecosystems in Australia: definitions and background*. In *Biodiversity of Mediterranean Ecosystems in Australia* (R. J. Hobbs, Ed, Surrey Beatty and Sons, Chipping Norton, New South Wales, Australia. pp 1-25.
- Hodgson, J. G., Thompson, K., Wilson, P. J., and Bogaard, A. (1998), Does biodiversity determine ecosystem function? The Ecotron experiment reconsidered. *Functional Ecology*. 12:843-856.
- Holben, B. (1986), Characteristics of maximum value composite images from temporal AVHRR data. *International Journal of Remote Sensing*. 7:1417-1434.
- Hooper, D. U. (1998), The role of complementarity and competition in ecosystem responses to variation in plant diversity. *Ecology*. 79(2):704-719.
- Hooper, D. U., and Vitousek, P. M. (1997), The effects of plant composition and diversity on ecosystem processes. *Science*. 277:1302-1305.
- Hooper, D. U., and Vitousek, P. M. (1998), Effects of plant composition and diversity on nutrient cycling. *Ecological Monographs*. 68(1):121-149.
- Huete, A. R. (1988), A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment*. 25:295-309.

- Hughes, J. B., and Petchey, O. L. (2001), Merging perspectives on biodiversity and ecosystem function. *Trends in Ecology and Evolution*. 16(5):222-223.
- Hurcom, S. J., and Harrison, A. R. (1998), The NDVI and spectral decomposition for semi-arid vegetation abundance estimation. *International Journal of Remote Sensing*. 19(16):3109-3125.
- Huston, M. A. (1997), Hidden treatments in ecological experiments: re-evaluating the ecosystem function of biodiversity. *Oecologia*. 110:449-460.
- Huston, M. A., Aarssen, L. w., Austin, M. P., Cade, B. S., Fridley, J. D., Garnier, E., Grime, J. P., Hodgson, J., Lauenroth, W. K., Thompson, K., Vandermeer, J. H., and Wardle, D. A. (2000), No consistent effect of plant diversity on productivity. *Science*. 289(25):1255a.
- Ikeda, H., Okamoto, K., and Fukuhara, M. (1999), Estimation of aboveground phytomass with a growth model using Landsat TM and climate data. *International Journal of Remote Sensing*. 20(11):2283-2294.
- Jasinski, M. F. (1990), Sensitivity of the normalized difference vegetation index to subpixel canopy cover, soil albedo, and pixel scale. *Remote Sensing of Environment*. 32:169-187.
- Johnson, K. H., Vogt, K. A., Clark, H. J., Schmitz, O. J., and Vogt, D. J. (1996), Biodiversity and the productivity and stability of ecosystems. *Trends in Ecology and Evolution*. 9(11):372-377.
- Jordan, C. F. (1969), Derivation of Leaf Area Index from quality of light on the forest floor. *Ecology*. 50:663-666.
- Journel, A. G., and Huilbregts, C. (1978), *Mining Geostatistics*. Academic Press, London, UK., pp. 600.
- Kabos, S., and Csillag, F. (Unpublished Manuscript), Simulating patchy landscapes with regional and local constraints.
- Karl, T. R., Heim, R. R., and Quayle, R. G. (1991), The greenhouse effect in central North America: if not now, when? *Science*. 251:1058-1061.
- Keddy. (1992), Assembly and response rules: two goals for predictive community ecology. *Journal of Vegetation Science*. NA(3):157-164.
- Keddy, P. A. (1990), *The use of functional as opposed to phylogenetic systematics: A first step in predictive community ecology*. In *Biological Approaches and Evolutionary trends in plants* (S. Kawano, Ed), Harcourt Brace Jovanovich, London, UK. pp 387-406.
- Kemp, and Williams. (1980), A physiological basis for niche separation between *agropyron smithii* (C3) and *bouteloua gracilis* (C4). *Ecology*. 4(61):846-858.
- Kephart, K. D., West, C. P., and Wedin, D. A. (1993), *Grassland ecosystems and their improvement*. In *Forages Volume I : An introduction to grassland agriculture* (R. F. Barnes, D. A. Miller, and C. J. Nelson, Eds.), Iowa State University Press, Ames, Iowa, USA. pp
- Kertész, M., and Csillag, F., (unpublished). HQSIM - HQ-Decomposition Algorithm.
- Knapp, A. K. (1993), Gas exchange dynamics in C3 and C4 grasses: consequences of differences in stomatal conductance. *Ecology*. 74(1):113-123.

- Knoop, W. T., and Walker, B. H. (1985), Interactions of woody and herbaceous vegetation in a southern African Savanna. *Journal of Ecology*. 83:235-253.
- Knops, J. M. H., Wedin, D., and Tilman, D. (2001), Biodiversity and decomposition in experimental grassland ecosystems. *Oecologia*. 126:429-433.
- Kolasa, J., and Rollo, C. D. (1991), *Heterogeneity of Heterogeneity*. In *Ecological Heterogeneity* (J. Kolasa, and S. T. A. Pickett, Eds.), Springer-Verlag, New York, NY, USA. pp1-23.
- Kutiel, P., and Danin, A. (1987), Annual-species diversity and above-ground phytomass in relation to some soil properties in the sand dunes of the northern Sharon Plains, Israel. *Vegetatio*. 70:45-59.
- Kumar, A. (1979), Assessment of net aerial primary production from harvest data using the "peak standing crop" and "difference" methods. *Comparative Physiological Ecology*. 4:62-63.
- Lachenbruch, P. A. (1975), *Discriminant Analysis*. . Hafner, New York, NY, USA, pp. 128.
- Larcher, W. (1995), *Physiological Plant Ecology: Ecophysiology and Stress Physiology and Functional Groups*. Springer-Verlag, Berlin, Germany. pp. 506.
- Lauenroth, W. K., Coffin, D. P., Burke, I. C., and Virginia, R. A. (1997), *Interactions between demographic and ecosystem processes in a semi-arid and arid grassland: a challenge for plant functional types*. In *Plant Functional Types. Their relevance to ecosystem properties and global change* (T. M. Smith, H. H. Shugart, and F. I. Woodward, Eds.), Cambridge University Press, Cambridge, UK. pp 234-254.
- Lauenroth, W. K., Sala, O. E., Coffin, D. P., and Kirchner, T. B. (1994), The importance of soil water in the recruitment of *bouteloua gracilis* in the shortgrass steppe. *Ecological Applications*. 4(4):741-749.
- Lauver, C. L., and Whistler, J. L. (1993), A hierarchical classification of Landsat TM imagery to identify natural grassland areas and rare species habitat. *Photogrammetric Engineering and Remote Sensing*. 59(5).
- Lawton, J. H. (2001), *Biodiversity, ecosystem processes and climate change*. In *Ecology: Achievement and Challenge* (M. C. Press, N. J. Huntly, and S. Levin, Eds.), Blackwell Science, London, UK. pp 139-160.
- Le Roux, X., and Bariac, T. (1998), Seasonal variations in soil, grass and shrub water status in a West African humid savanna. *Oecologia*. 113:456-466.
- Legendre, P., and Fortin, M.-J. (1989), Spatial pattern and ecological analysis. *Vegetatio*. 80:107-138.
- Lehman, C. L., and Tilman, D. (2000), Biodiversity, stability, and productivity in competitive communities. *The American Naturalist*. 165(5):534-552.
- Leonard, R. D., and Dobson, R. B. (1991), *Park Conservation Plan. Grasslands National Park*. Prairie and Northern Region, Canadian Parks Service.
- Leps, J., Osbornová-Kosinová, M., and Rejmaninek, M. (1982), Community stability, complexity and species life-history strategies. *Vegetatio*. 50:53-63.
- Levin, S. A. (1992), The problem of pattern and scales in ecology. *Ecology*. 73:1943-1967.

- Levin, S. A. (1993), *Concepts of scale at the local level*. In *Scaling physiological processes. Leaf to Globe* (J. E. Ehleringer, and C. B. Field, Eds.), Academic Press, New York, NY, USA.
- Lewis, W. M. J. (1978), Comparison of temporal and spatial variation in the zooplankton of a lake by means of variance components. *Ecology*. 59:666-671.
- Lobo, A., Moloney, K., Chic, O., and Chiariello, N. (1998), Analysis of fine-scale spatial pattern of a grassland from remotely-sensed imagery and field collected data. *Landscape Ecology*. (13):111-131.
- Loreau, M. (1998a), Biodiversity and ecosystem functioning: A mechanistic model. *Proceedings of the National Academy of Sciences*. 95:5632-5636.
- Loreau, M. (1998b), Separating sampling and other effects in biodiversity experiments. *OIKOS*. 82:600-602.
- Loreau, M., and Hector, A. (2001), Partitioning selection and complementarity in biodiversity experiments. *Nature*. 412:72-76.
- Loreau, M., Naeem, S., Inchausti, P., Bentgsson, J., Grime, J. P., Hector, A., Hooper, D. U., Huston, M. A., Raffaelli, D., Schmid, B., Tilman, D., and Wardle, D. A. (2001), Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science*. 294:804-808.
- Loveridge, D. M., and Potyondi, B. (1983), *From Wood Mountain to the Whitemud: A historical survey of the Grasslands National Park area*. History and Archaeology 67, National Historic Parks and Sites Branch, Parks Canada, Environment Canada, Ottawa.
- Ludwig, J. A., and Goodall, D. W. (1978), A comparison of paired- with blocked-quadrat variance methods for the analysis of spatial pattern. *Vegetatio*. 38:49-59.
- MacGillivray, G. W., Grime, J. P., Band, S. R., Boothe, R. E., Campbell, B., Hendry, G. A. F., Hillier, S. H., Hodgson, J. G., Hunt, R., Jalili, A., Mackey, J. M. L., Mowforth, M. A., Neal, A. M., Reader, R., Rorison, H. I., Spencer, R. E., Thompson, K., and Thorpe, P. C. (1995), Testing predictions of the resistance and resilience of vegetation subject to extreme events. *Functional Ecology*. 9(4):640-649.
- Madson, J. (1997), *Tallgrass prairie*. Falcon Publishing, pp. 321.
- Magurran, A. E. (1988), *Ecological diversity and its measurement*. . Croom Helm, London, UK, pp. 179.
- Malingreau, J. P. (1986), Global vegetation in dynamics, satellite observations over Asia. *International Journal of Remote Sensing*. 7:1121-1146.
- Malone, C. R. (1968), Determination of peak standing crop biomass of herbaceous shoots by the harvest method. *American Midland Naturalist*. 79:429-435.
- McBratney, A. B., and Webster, R. (1986), Choosing functions for semi-variograms of soil properties and fitting them to sampling estimates. *Journal of Soil Science*. 37:617-639.

- McCanny, S. (2000), *Guide to the plants of Grasslands National Park*. In *Grasslands National Park Field Guide* (K. S. Fargey, S. D. Larson, S. J. Grant, P. Fargey, and C. Schmidt, Eds.), Prairie Wind and Silver Sage - Friends of Grasslands Inc., Swift Current, Sk, Canada.
- McCloy, K. R., Schoneveld, R., and Kemp, D. (1993), Measurement of pasture parameters from reflectance data. *International Journal of Remote Sensing*. :1107-1118.
- McIntosh, R. P. (1991), *Concept and terminology of heterogeneity in ecology*. In *Ecological Heterogeneity* (J. Kolasa, and S. T. A. Pickett, Eds.), Springer-Verlag, New York, NY, USA. pp 24-46.
- McLachlan, G. J. (1992), *Discriminant Analysis and Statistical Pattern Recognition*. . John Wiley and Sons, New York, NY, USA. pp. 544.
- McNaughton, S. J. (1977), Diversity and stability of ecological communities: a comment on the role of empiricism in ecology. *American Naturalist*. 111:515-525.
- Menard, S. (1995), *Applied logistic regression analysis*. Quantitative Applications in the Social Sciences No. 106. Sage Publications, Thousand Oakes, CA, USA, pp. 98.
- Michalsky, S. J., and Ellis, R. A. (1994), *Vegetation of Grasslands National Park*. DA Westworth and Associates, Calgary.
- Miesch, A. T. (1975), Variograms and variance components in geochemistry and ore evaluation. *Geological Society of America Memoir*. 142:333-340.
- Milner, C., and Hughes, R. E., (1968), *Methods for the measurement of primary production of grassland*. IBP Handbook, No. 6, Blackwell, Oxford, UK.
- Mino, N., Saito, G., and Ogawa, S. (1998), Satellite monitoring of changes in improved grassland management. *International Journal of Remote Sensing*. 19(3):439-452.
- Mitchell, S. W., and Csillag, F. (2001), Assessing the stability and uncertainty of vegetation growth under climatic variability: northern mixed grass prairie. *Ecological Modelling*. 139(2-3):101-121.
- Monson, R. K., Edwards, G. E., and Ku, M. S. B. (1984), C3-C4 intermediate photosynthesis in plants. *BioScience*. 34(9):563-574.
- Mooney, H. A., Cushman, J. H., Medina, E., Sala, O. E., and Schulze, E. D. (1996), *Functional roles of biodiversity: a global perspective*. John Wiley and Sons, Chichester, UK, pp. 493.
- Mooney, H. A., Lubchenco, J., Dirzo, R., and Sala, O. E. (1995), *Biodiversity and ecosystem functioning: basic principles*. In *Global Biodiversity Assessment* (V. H. Heywood, and R. T. Watson, Eds.), Cambridge University Press, Cambridge, UK. pp 275-326.
- Myeni, R. B., Ross, J., and Asrar, G. (1989), A review on the theory of photon transport in leaf canopies. *Agricultural and Forest Meteorology*. 45:1-153.
- Naeem, S. (2000), Reply to Wardle et al. *Bulletin of the Ecological Society of America*. 81:241-246.
- Naeem, S., Håkansson, K., Lawton, J. H., Crawley, M. J., and Thompson, L. J. (1996), Biodiversity and plant productivity in a model assemblage of plant species. *OIKOS*. 76:259-264.

- Naeem, S., and Li, S. (1997), Biodiversity enhances ecosystem reliability. *Nature*. 390:507-509.
- Naeem, S., Thompson, L. J., Lawler, S. P., Lawton, J. H., and Woodfin, R. M. (1994), Declining biodiversity can alter the performance of ecosystems. *Nature*. 368:734-737.
- Naeem, S., Thomson, L. J., Lawlor, S., Lawton, J. H., and Woodfin, R. M. (1995), Empirical evidence that declining species diversity may alter the performance of terrestrial ecosystems. *Philosophical Transactions of the Royal Society of London*. 347:249-262.
- Neilson, R. P. (1993), Transient ecotone response to climatic change: some conceptual and modelling approaches. *Ecological Applications*. 3(3):385-395.
- Nelder, J. A., and Wedderburn, R. W. M. (1972), Generalized linear models. *Journal of the Royal Statistical Society*. 135:370-384.
- Noble, I. R. (1989), *Attributes of invaders and the invading process: terrestrial and vascular plants*. In *Biological invasions : a Global Perspective* (J. A. Drake, H. A. Mooney, F. di Castri, R. H. Groves, F. J. Kruger, M. Rejmanek, and M. Williamson, Eds.), John Wiley & Sons Ltd, Chichester, UK. pp 301-311.
- Nortcliff, S. (1978), Soil variability and reconnaissance soil mapping: a statistical study in Norfolk. *Journal of Soil Science*. 29:403-418.
- Noy-Meir, I. (1973), Desert ecosystems: environment and producers. *Annual Review of Ecological Systems*. 4:25-51.
- Ode, D., and Tieszen, L. L. (1980), The seasonal contribution of C3 and C4 plant species to primary production in a mixed prairie. *Ecology*. 61:1304-1311.
- Ojima, D. S., Parton, W. J., Coughenor, M. B., Scurlock, J. M. O., Kirchner, T. B., Kittel, T. G. F., Hall, D. O., Schimel, D. S., Moya, E. G., Gilmarov, T. G., Seastedt, T. R., Kamnalrut, A., Kinyamario, J. I., Long, S. P., Menaut, J.-C., Sala, O. E., Scholes, R. J., and Veen, J. A. (1996), *Impact of climate and atmospheric carbon dioxide changes on grasslands of the world*. In *Global Change: Effects on Coniferous Forests and Grasslands* (D. O. H. Breymer, J. M. Melillo, and G. I. Ågren, Eds.), John Wiley and Sons. pp 273-311.
- Olden, J. D., and Jackson, D. A. (2000), Torturing data for the sake of generality: How valid are our regression models? *EcoScience*. 7(4):501-510.
- Oliver, M. A., and Webster, R. (1986), Combining nested and linear sampling for determining the scale and form of spatial variation of regionalized variables. *Geographical Analysis*. 18:227-242.
- O'Neill, (1996), Scale problems in reporting landscape pattern at the regional scale. *Landscape Ecology*. 3(11):169-180.
- O'Neill, R. V., R. G., Milne, B. T., Turner, M. G., and Jackson, B. (1991), *Heterogeneity and spatial hierarchies*. In *Ecological Heterogeneity* (J. Kolasa, and S. T. A. Pickett, Eds.), Springer-Verlag, New York, NY, USA. pp85-96.

- Ovington, J. D., Heitcamp, D., and Lawrence, D. B. (1963), Plant biomass and productivity of prairie savanna, oakwood and maize field ecosystems in central Minnesota. *Ecology*. 44:52-65.
- Owensby, C. E., Coyne, P. I., Ham, J. M., Auen, L. A., and Knapp, A. K. (1993), Biomass production in a tallgrass prairie ecosystem exposed to ambient and elevated CO₂. *Ecological Applications*. 3:664-653.
- Paine, R. T. (1980), Food webs: linkage interaction strength and community infrastructure. *Journal of Animal Ecology*. 49:667-685.
- Paine, R. T., and Levin, S. A. (1981), Intertidal landscapes: disturbance and the dynamics of patches. *Ecological Monographs*. 51:145-178.
- Parton, W. J., Coughenor, M. B., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner, T. B., Menaut, J.-C., Seastedt, T. R., Moya, E. G., Kamnalrut, A., Kinyamario, J. I., and Hall, D. O. (1996), *Global grassland ecosystem modelling: development and test of ecosystem models for grassland systems*. In *Global Change: Effects on Coniferous Forests and Grasslands* (D. O. Breymer, J. M. Melillo, and G. I. Ågren, Eds.), John Wiley and Sons. pp 229-269.
- Paruelo, J. M., and Lauenroth, w. k. (1995), Regional patterns of normalized difference vegetation index in North American shrublands and grasslands. *Ecology*. 76(6):1888-1898.
- Pearcy, R. W., and Ehleringer, J. (1984), Comparative ecophysiology of C₃ and C₄ plants. *Plant, Cell and Environment*. 7:1-13.
- Peat, H. C. L. (1997), *Dynamics of C₃ and C₄ productivity in northern mixed grass prairie*, M.Sc Thesis, University of Toronto, Toronto, pp. 163.
- Pickett, S. T. A., and Cadenasso, M. L. (1995), Landscape ecology: spatial heterogeneity in ecological systems. *Science*. 269:331-334.
- Pickup, G., Chewings, V. H., and Nelson, D. J. (1993), Estimating changes in vegetation cover over time in arid rangelands using landsat MSS data. *Remote Sensing of Environment*. 43:243-263.
- Piper, J. K. (1995), Composition of prairie plant communities on productive versus unproductive sites in wet and dry years. *Canadian Journal of Botany*. 73:1635-1644.
- Poole, D. K., and Miller, P. C. (1975), Water relations of selected species of chaparral and coastal sage communities. *Ecology*. 56:1118-1128.
- Puerto, A., Rico, M., Matias, M. D., and Garcia, J. A. (1990), Variation in structure and diversity in Mediterranean grasslands related to trophic status and grazing intensity. *Journal of Vegetation Science*. 1:445-452.
- Purevdorj, T., Tateishi, R., Ishiyama, T., and Honda, Y. (1998), Relationships between percent vegetation cover and vegetation indices. *International Journal of Remote Sensing*. 19(18):3519-3535.
- Purvis, A., and Hector, A. (2000), Getting the measure of biodiversity. *Nature*. 405(11 May 2000):212-219.

- Qi, J., Chehbouni, A., Huete, A. R., Kerr, Y. H., and Sorooshian, S. (1994), A modified soil adjusted vegetation index. *Remote Sensing of Environment*. 48:119-126.
- Quattrochi, D., and Pelletier, R. E. (1991), *Remote sensing of landscapes : an introduction*. In *Quantitative Methods in Landscape Ecology* (M. Turner, and R. H. Gardner, Eds.), Springer Verlag, New York. pp51-76.
- Reed, B. C., Brown, J. F., VanderZee, D., Loveland, T. R., Merchant, J. W., and Ohlen, D. O. (1994), Measuring phenological variability from satellite imagery. *Journal of Vegetation Science*. 5:703-714.
- Richardson, A. J., and Everitt, J. H. (1992), Using spectral vegetation indices to estimate rangeland productivity. *Geocarto International*. 1:63-69.
- Richardson, A. J., and Wiegand, C. L. (1977), Distinguishing vegetation from soil background information. *Photogrammetric Engineering and Remote Sensing*. 43:1541-1552.
- Ricklefs, R. E. (1990), *Ecology*. . Freeman, New York, NY, USA, pp. 896.
- Riera, J. L., Magnuson, J. J., Vande Castle, J. R., and MacKenzie, M. D. (1998), Analysis of large-scale spatial heterogeneity in vegetation indices among North American landscapes. *Ecosystems*. 1:268-282.
- Ripley, B. D. (1996), *Pattern Recognition and Neural Networks*. Cambridge University Press, Cambridge, UK, pp. 403.
- Risser, P. G. (1985), *Grasslands*. In *Physiological Plant Ecology of North American Plant Communities* (B. F. Chabot, and H. A. Mooney, Eds.). pp 233-256.
- Ross, J. K. (1981), *The radiation regime and architecture of plant stands*. Kluwer Boston, Hingham, MA, USA, pp. 480.
- Rossi, R. E., Mulla, D. J., Journel, A. G., and Franz, E. H. (1992), Geostatistical tools for modeling and interpreting ecological spatial dependence. *Ecological Monographs*. 62:277-314.
- Roujean, J.-L., and Breon, F. M. (1995), Estimating PAR absorbed by vegetation from bidirectional reflectance measurements. *Remote Sensing of Environment*. 51:375-384.
- Rouse, J. W., Haas, R. H., Schell, J. A., and Deering, D. W., 1973. Monitoring vegetation systems in the Great Plains with ERTS-1, 3rd Earth Resources Technology Satellite Symposium, pp. 309-317.
- Rowe, J. S. (1980), *Landscapes: A guide to landforms and ecology of southern Saskatchewan*. (R. J. Long, Ed). Saskatchewan Environment, Regina, Saskatchewan, Canada. pp101.
- Roy, P. S., Jonna, S., and Pant, D. N. (1991), Evaluation of grasslands and spectral reflectance relationship to its biomass in Kanha National Park (M.P.), India. *Geocarto International*. 1:39-45.
- Running, S. W., Nemani, R. R., Peterson, D. L., Band, L. E., Potts, D. F., Pierce, L. L., and Spanner, M. A. (1989), Mapping regional evapotranspiration and photosynthesis by coupling satellite data with ecosystem simulation. *Ecology*. 70:1090-1101.

- Sage, R F, Wedin, D A, Li, M. (1999). *The biogeography of C4 photosynthesis: Patterns and controlling factors*. In *C4 Plant Biology* (R.F Sage and K. Monson, Eds.), Academic Press, San Diego, CA, USA. pp313-356.
- Sala, O. E., Golluscio, R. A., Lauenroth, W. K., and Soriano, A. (1989), Resource partitioning between shrubs and grasses in the Patagonian Steppe. *Oecologia*. 81:501-505.
- Sala, O. E., Lauenroth, W. K., and Golluscio, R. A. (1997), *Plant functional types in temperate semi-arid regions*. In *Plant Functional Types. Their relevance to ecosystem properties and global change* (T. M. Smith, H. H. Shugart, and F. I. Woodward, Eds.), Cambridge University Press, Cambridge, UK. pp 217-233.
- Samson, F. B., and Knopf, F. L. (1994), Prairie conservation in North America. *BioScience*. 44:418-421.
- Samson, F. B., Knopf, F. L., and Ostlie, W. R. (1998), *Grasslands*. In *Status and Trends of the Nation's Biological Resources* (M. J. Mac, P. A. Opler, C. E. Puckett, and P. D. Doran, Eds.), U.S Department of the Interior, U.S. Geological Survey, Reston, VA, USA. pp437-472.
- Saskatchewan Institute of Pedology, (1992), *Grasslands National Park Soil Survey*. University of Saskatchewan, Saskatoon.
- Scheffé, H. (1959), *The analysis of variance*. John Wiley and Sons, New York, NY, USA, pp. 477.
- Schläpfer, F., and Schmid, B. (1999), Ecosystem effects of biodiversity: a classification hypothesis and exploration of empirical results. *Ecological Applications*. 9:893-912.
- Schläpfer, F., Schmid, B., and Seidl, I. (1999), Expert estimates about effects of biodiversity on ecosystem processes and services. *OIKOS*. 84:346-352.
- Scholes, R. J., Pickett, G., Ellery, W. N., and Blackmore, A. C. (1997), *Plant functional types in African savannas and grasslands*. In *Plant Functional Types* (T. M. Smith, H. H. Shugart, and F. I. Woodward, Eds.), Cambridge University Press, Cambridge. pp255-270.
- Schulze, E.-D., Mooney, H. A., Sala, O. E., Jobbagy, E., Buchmann, N., Bauer, G., Canadell, J., Jackson, R. B., Loreti, J., Oesterheld, M., and Ehleringer, J. R. (1996), Rooting depth, water availability, and vegetation cover along an aridity gradient in Patagonia. *Oecologia*. 108:53-511.
- Schuster, W. S., and Monson, R. K. (1990), An examination of the advantages of C3-C4 intermediate photosynthesis in warm environments. *Plant, Cell and Environment*. 13:903-912.
- Schwartz, M. W., Brigham, C. A., Hoeksema, J. D., Lyons, K. G., Mills, M. H., and van Mantgem, P. J. (2000), Linking biodiversity to ecosystem function: Implications for conservation ecology. *Oecologia*. 122:297-305.
- Seastedt, T. R., Briggs, J. M., and Gobson, D. J. (1991), Controls of nitrogen limitation in tallgrass prairie. *Oecologia*. 87:72-79.
- Sellers, P. J., Hall, F. G., Strebel, D. H., Asrar, G., and Murphy, R. E. (1990), *Satellite remote sensing and field experiments*. In *Remote Sensing of Biosphere Functioning* (R. J. Hobbs, and H. A. Mooney, Eds.), Springer-Verlag, New York. pp169-202.

- Senft, R. L., Coughenour, M. B., Bailey, D. W., Rittenhouse, L. R., Sala, O. E., and Swift, D. E. (1987), Large herbivore foraging and ecological hierarchies. *BioScience*. 37:789-799.
- Shao, J., and Tu, D. (1995), *The Jackknife and Bootstrap*. Springer-Verlag, New York, NY, USA, pp. 536.
- Shmida, A., and Ellner, S. (1989), Coexistence of plant species with similar niches. *Vegetatio*. 58:29-55.
- Simberloff, D., and Dayan, T. (1991), The guild concept and the structure of ecological communities. *Annual Review of Ecology and Systematics*. 22:115-143.
- Singh, J. S., Lauenroth, W. K., and Milchunas, D. G. (1983), Geography of grassland ecosystems. *Progress in Physical Geography*. 7:46-80.
- Smith, T., and Huston, M. (1989), A theory of the temporal and spatial dynamics in plant communities. *Vegetatio*. 83:49-70.
- Sokal, R. R., and Rohlf, F. J. (1981), *Biometry. The Principles and Practice of Statistics in Biological Research*. W.H. Freeman and Company, New York, pp. 859.
- Soriano, A., and Sala, O. E. (1983), Ecological strategies in a Patagonian arid steppe. *Vegetatio*. 56:9-15.
- Spehn, E. M., Joshi, J., Schmid, B., Diemer, M., and Körner, C. (2000), Above-ground resource use increases with plant species richness in experimental grassland ecosystems. *Functional Ecology*. 14:326-337.
- Stoms, D. M., and Estes, J. E. (1993), A remote sensing research agenda for mapping and monitoring biodiversity. *International Journal of Remote Sensing*. 20(10):1839-1860.
- Stone, M. (1974), Cross-validatory choice and assessment of statistical predictions. *Journal of the Royal Statistical Society*. 36:111-133.
- Stubbendieck, K., Hatch, S. L., and Butterfield, C. H. (1992), *North American Range Plants*. University of Nebraska Press, NE, USA. pp 493.
- Swaine, M. D., and Whitmore, T. C. (1988), On the definition of ecological species groups in tropical rain forests. *Vegetatio*. 75:81-86.
- Swift, M. J., and Anderson, J. M. (1993), *Biodiversity and ecosystem function in agricultural systems*. In *Biodiversity and Ecosystem Function* (E.-D. Schulze, and H. A. Mooney, Eds.), Springer-Verlag, Berlin, Germany. pp15-42.
- Symstad, A. J. (2000), A test of the effects of functional group richness and composition on grassland invasability. *Ecology*. 81(1):99-109.
- Symstad, A. J., Tilman, D., Wilson, J., and Knops, J. M. H. (1998), Species loss and ecosystem functioning: effects of species identity and community composition. *OIKOS*. 81:389-397.
- Szaro, R. C. (1986), Guild management: an evaluation of avian guilds. *Environmental management*. 10:681-688.
- Teeri, J. A., and Stowe, K. G. (1976), Climate patterns and distribution of C4 grasses in North America. *Oecologia*. 23:1-12.

- Teese. (1995), Intraspecific variation for CO₂ compensation point and differential growth among variants in a C3-C4 intermediate plant. *Oecologia*. NA(102):371-376.
- Tieszen, L. L. (1970), Photosynthetic properties of some grasses in eastern South Dakota. *Proceeds of the Dakota Academy of Science*. 49:78-89.
- Tieszen, L. L., Reed, B. C., Bliss, N. B., Wylie, B. K., and DeJong, D. D. (1997), NDVI, C3 and C4 production, and distributions in Great Plains grassland land cover classes. *Ecological Applications*. 7(1):59-78.
- Tieszen, L. L., Senyimba, M. M., Imbamba, S. K., and Troughton, J. H. (1979), The distribution of C3 and C4 grasses and carbon isotope discrimination along an altitudinal and moisture gradient in Kenya. *Oecologia*. 37:377-350.
- Tilman, D. (1999), The ecological consequences of biodiversity: a search for general principles. *Ecology*. 80(5):1455-1474.
- Tilman, D. (2000), Causes, consequences and ethics of biodiversity. *Nature*. 405(11 May 2000):208-211.
- Tilman, D. (2001), *Effects of diversity and composition on grassland stability and productivity*. In *Ecology: Achievement and Challenge* (M. C. Press, N. J. Huntly, and S. A. Levin, Eds.), Blackwell Science, London, UK. pp183-207.
- Tilman, D., and Downing, J. A. (1994), Biodiversity and stability in grasslands. *Nature*. (367):363-365.
- Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M., and Siemann, E. (1997a), The influence of functional diversity and composition on ecosystem processes. *Science*. 277:1300-1302.
- Tilman, D., Lehman, C. L., and Thomson, K. T. (1997b), Plant diversity and ecosystem productivity: theoretical considerations. *Proceedings of the Natural Academy of Sciences*. 94:1857-1861.
- Tilman, D., Reich, P. B., Knops, J., Wedin, D., Mielke, T., and Lehman, C. (2001), Diversity and productivity in a long-term grassland experiment. *Science*. 294:843-845.
- Tilman, D., Wedin, D., and Knops, J. (1996), Productivity and sustainability influenced by diversity in grassland ecosystems. *Nature*. 379:718-720.
- Todd, S. W., Hoffer, R. M., and Milchunas, D. G. (1998), Biomass estimation on grazed and ungrazed rangelands using spectral indices. *International Journal of Remote Sensing*. 19(3):427-438.
- Tòth, T., Csillag, F., Biehl, L. L., and Micheli, E. (1991), Characterization of semivegetated salt-affected soils by means of field remote sensing. *Remote Sensing of Environment*. 37:167-180.
- Tucker, C. J. (1979), Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment*. 8:127-150.
- Tucker, C. J. (1980), Remote sensing of leaf water content in the near infrared. *Remote Sensing of Environment*. 10:23-32.
- Tucker, C. J., Newcomb, W. W., Los, S. O., and Prince, S. D. (1991), Mean and inter-year variation of growing-season normalized difference vegetation index for the Sahel 1981-1989. *International Journal of Remote Sensing*. 12(6):1133-1135.

- Tucker, C. J., Vanpraet, C. L., Sharman, M. J., and Van Ittersum, G. (1983), Satellite remote sensing of total herbaceous biomass in the Senegalese Sahel: 1980-1984. *Remote Sensing of Environment*. 17:233-249.
- Turner, C. L., Seastedt, T. R., Dyer, M. I., Kittel, G. F., and Schimel, D. S. (1992), Effects of management and topography on the radiometric response of a tallgrass prairie. *Journal of Geophysical Research*. 97(D17):18855-18866.
- Turner, M. G. (1989a), Landscape ecology: the effect of pattern on process. *Annual Review of Ecology and Systematics*. 20:171-197.
- Turner, M. G. (1989b), *Landscape heterogeneity and disturbance*. Springer-Verlag, New York, NY, USA, pp. 239.
- Urban, D. L., O'Neill, R. V., and Shugart, H. H. J. (1987), Landscape ecology. *Bioscience*. 37(119-127).
- Valentine, J. F. (1990), *Grazing Management*. Academic Press, New York, NY, USA, pp. 533.
- van der Hiejen, M. G. A., Klironomos, J. N., Ursic, M., Moutoglou, P., Streitwolf-Engel, R., Boller, T., Wiemken, A., and Sanders, I. R. (1998), Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*. 369:69-72.
- Venables, W. N., and Ripley, B. D. (1997), *Modern applied statistics with S-PLUS*. Statistics and computing. Springer-Verlag, New York, pp. 501.
- Ver Hoef, J. M., Cressie, N. A. C., and Glenn-Lewin, D. C. (1993), Spatial models for spatial statistics: some unification. *Journal of Vegetation Science*. 4:441-452.
- Vermote, E. F., Tanré, D., Deuze, J. L., Herman, M., and Morcrette, J.-M. (1997), Second simulation of the satellite signal in the solar spectrum, 6s: An overview. *IEEE Transactions on Geoscience and Remote Sensing*. 35(3):675-686.
- Verner, J. (1984), The guild concept applied to management of bird populations. *Environmental Management*. 8:1-14.
- Vestergaard, P. M., and Hooper, D. U. (1994), Response to mowing of coastal brackish meadow plant communities along an elevational gradient. *Nordic Journal of Botany*. 14:569-587.
- Vitousek, P. M., and Hooper, D. U. (1993), *Biological diversity and terrestrial ecosystem biogeochemistry*. In *Biodiversity and Ecosystem Function* (E.-D. Schulze, and H. A. Mooney, Eds.), Springer-Verlag, Berlin, Germany. pp 3-14.
- Vitousek, P. M., Mooney, H. A., Lubchenko, J., and Melillo, J. M. (1997), Human domination of the Earth's ecosystems. *Science*. 277:494-499.
- Vitousek, P. M., Walker, R. M., Whiteaker, L. D., Mueller-Dombois, P. A., and Matson, A. (1987), *Science*. 238:802.
- Waide, R. B., Willig, M. R., Steiner, C. F., Mittelbach, G., Gough, L., Dodson, S. I., Juday, J. P., and Parmenter, R. (1999). The relationship between productivity and species richness. *Annual Review of Ecology and Systematics*. 30: 257-300.

- Walker, B. H. (1992), Biodiversity and ecological redundancy. *Conservation Biology*. 6:18-23.
- Walsh, S. J., Moody, A., Allen, T. R., and Brown, D. G. (1997), *Scale dependence of NDVI and its relationship to mountainous terrain*. In *Scale in Remote Sensing and GIS* (D. A. Quattrochi, and M. F. Goodchild, Eds.), CRC Press, Boca Raton, FL, USA. pp27-56.
- Walter, H. (1979), *Vegetation of the earth*. . Springer, New York, USA., pp. 274.
- Walthall, C. L., and Middleton, E. M. (1992), Assessing spatial and seasonal variations in grasslands with spectral reflectances from a helicopter platform. *Journal of Geophysical Research*. 97:18905-18912.
- Wardle, D. A., Bonner, K. I., and Nicholson, K. S. (1997a), Biodiversity and plant litter: experimental evidence which does not support the view that enhanced species richness improves ecosystem function. *OIKOS*. 79:247-258.
- Wardle, D. A., Zackrisson, O., Hörnberg, G., and Gallet, C. (1997b), The influence of island area on ecosystem properties. *Science*. 277:1296-1299.
- Webster, R. (1978), Optimally partitioning soil transects. *Journal of Soil Science*. 29:388-402.
- Webster, R. (1979), *Nested Classification and Sampling*. In *Quantitative numerical methods in soil classification survey* (R. Webster, Ed), Oxford University Press, Bristol, UK. pp 90-106.
- Webster, R., Curran, P. J., and Munden, J. W. (1989), Spatial correlation in reflected radiation from the ground and its implications for sampling and mapping by ground-based radiometry. *Remote Sensing of Environment*. 29:67-78.
- Webster, R., and Oliver, M. A. (1990a), *Nested Sampling and Analysis*. In *Statistical Methods in Soil and Land Resource Survey* (R. Webster, and M. A. Oliver, Eds.), Oxford University Press, New York, USA. pp241-259.
- Webster, R., and Oliver, M. A. (1990b), *Spatial Dependence*. In *Statistical Methods in Soil and Land Resource Survey* (R. Webster, and M. A. Oliver, Eds.), Oxford University Press, New York, USA. pp214-240.
- Webster, R., and Oliver, M. A. (1992), Sampling adequately to estimate variograms of soil properties. *Journal of Soil Science*. 43:177-192.
- Weiser, R. L., Asrar, G., Miller, G. P., and Kanemasu, E. T. (1986), Assessing grassland biophysical characteristics from spectral measurements. *Remote Sensing of Environment*. 20:141-152.
- Weltzin, J. F., and McPherson, G. R. (1997), Spatial and temporal soil moisture resource partitioning by trees and grasses in a temperate savanna, Arizona, USA. *Oecologia*. 112:156-164.
- Westoby, M., and Leishman, M. (1997), *Categorizing plant species into functional types*. In *Plant Functional Types. Their relevance to ecosystem properties and global change* (T. M. Smith, H. H. Shugart, and F. I. Woodward, Eds.), Cambridge University Press, Cambridge, UK. pp104-121.
- White, R., Murray, S., and Rohweder, M. (2000), *Pilot Analysis of Global Ecosystems: Grassland Ecosystems*. World Resources Institute, Washington, DS, USA. pp80.

- Whittaker, R. H. (1975), *Communities and Ecosystems*. . Macmillan Publishing, New York, USA, pp. 385 .
- Whittaker, R. H., and Likens, G. E. (1973), *Carbon in the biota*. In *Carbon and Biosphere* (G. M. Woodwell, and E. V. Pecan, Eds.), National Technical Information Service, Springfield. pp281-302.
- Wickham, J. D., and Riitters, K. H. (1995), Sensitivity of landscape metrics to pixel size. *International Journal of Remote Sensing*. 16(18):3585-3594.
- Wilsey, B. J., and Potvin, C. (2000), Biodiversity and ecosystem functioning: Importance of species evenness in an old field. *Ecology*. 81(4):887-892.
- Wu, J., and Levin, S. A. (1994), A spatial patch dynamic modeling approach to pattern and process in an annual grassland. *Ecological Monographs*. 4(64):447-464.
- Wu, J., and Levin, S. A. (1997), A patch-based spatial modeling approach: conceptual framework and simulation scheme. *Ecological Modelling*. 101:325-346.
- Yang, L., Wylie, B. K., Tieszen, L. L., and Reed, B. C. (1998), An analysis of relationships among climate forcing and time-integrated NDVI of grasslands over the U.S. Northern and Central Great Plains. *Remote Sensing of Environment*. 65:25-37.
- Yodzis, P. (1982), The compartmentation of real and assembled ecosystems. *American Naturalist*. 120:551-570.
- Zalom, F. G., Goodell, P. B., Wilson, L. T., Barnett, W. W., and Bentley, W. J. (1983), *Degree-days: The calculation and use of heat units in pest management*. Leaflet 21373, Division of Agriculture and Natural Resources, University of California.
- Zar, J. H. (1984), *Biostatistical Analysis*. Prentice-Hall, Inc., Englewood Cliffs, N.J., USA, pp. 718.
- Zimmerman, D. L., and Zimmerman, M. B. (1991), A comparison of spatial semivariogram estimators and corresponding ordinary kriging predictors. *Technometrics*. 33:77-92.